### fieldbean

# **CAPRI**


## High yield with high protein content



#### **Benefits:**

- Very high seed yields with high protein yield per ha
- Uniform seed size (500-600 g TGW)
- Early flowering and maturity
- Good stiffness of straw
- Easy to combine
- High feed value

#### Variety characteristics: (Officially confirmed or respectively in line with Bundessortenamt)



Breeder: P. H. PETERSEN Saatzucht Lundsgaard GmbH, Version: 09.01.2023 / 18.00







## High yield with high protein content

Us	age:			
_				

Grain use Nitrogen fixation Suitable for human and animal nutrition

#### **Cultivation recommendations:**

Recommended sowing rate	40 - 50 seeds/m²			
Sowing depth	6 - 8 cm on heavy soils, 8 - 10 cm on light soils			
Sowing period	Earliest possible in spring - frost tolerant up to approx5°C			
Sowing method	Single or double row spacing is beneficial for the yield, larger row spacing in combination with precision sowing is possible.			
Fertilization	No N-fertilization necessary. Depending on the nutrient content of the soil and nutrient extraction 40-60 kg $P_2O_5$ /ha, 100-130 kg $K_2O$ /ha and about 20-50 kg MgO/ha are advised. If necessary an application of sulphur is suggested.			
Crop protection	Use herbicides during pre-emergence or mechanically hoe or harrow the field for weed control. Post-emergence herbicide application specifically against grasses is possible. If necessary use fungizides against chocolate spots. Use clean seed to avoid ascochyta blight. When considering usage of insecticides against black bean aphid or bean weevil mind the damage threshhold.			
Harvest	Easy to combine due to even maturity and high stiffness. Optimal moister for the harvest is 15-19%. Use protective setting for threshing to avoid broken grains.			



